




ASK YOUR DENTIST ABOUT

EMERGENCY CARE



AMERICAN ACADEMY OF PEDIATRIC DENTISTRY



**When your child needs urgent dental treatment,
your pediatric dentist stands ready to help.
Please keep the emergency number
available and convenient.**

EMERGENCY

What should I do if my child's baby tooth is knocked out?

Contact your pediatric dentist as soon as possible. The baby tooth should not be replanted because of the potential for subsequent damage to the developing permanent tooth.

What should I do if my child's permanent tooth is knocked out?

Find the tooth and rinse it gently in cool water. (Do not scrub or clean it with soap — use only water!) If possible, replace the tooth in the socket immediately and hold it there with clean gauze or a wash cloth. If you can't put the tooth back in the socket, place the tooth in a clean container with cold milk, saliva or water. Get to the pediatric dental office immediately. (Call the emergency number if it's after hours.) The faster you act, the better your chances of saving the tooth.

What if a tooth is chipped or fractured?

Contact your pediatric dentist immediately. Quick action can save the tooth, prevent infection and reduce the need for extensive dental treatment. Rinse the mouth with water and apply cold compresses to reduce swelling if the lip also was injured. If you can find the broken tooth fragment, place it in cold milk or water and bring it with you to the dental office.



What about a severe blow to the head or a jaw fracture?

You need immediate medical attention. A severe head injury can be life threatening. Keep in mind that an emergency medical team might be able to reach you faster than you can get to the hospital.

CARE

What if my child has a toothache?

Call your pediatric dentist and visit the office promptly. To comfort your child, rinse the mouth with water. Over-the-counter children's pain medication, dosed according to your child's weight and age, might ease the symptoms. You may apply a cold compress or ice wrapped in a cloth to the face in the area of the pain, but do not put heat or aspirin on the sore area.



Can dental injuries be prevented?

Your child's risk for dental injuries can be reduced greatly by following a few simple suggestions. First, reduce risk for severe oral injury in sports by wearing protective gear, including a mouthguard. Second, always use a car seat for young children and require seat belts for everyone else in the car. Third, child-proof your home to prevent falls and electrical injuries. Regular dental check-ups provide your dentist an opportunity to discuss additional age-appropriate preventive strategies with your child.

The American Academy of Pediatric Dentistry (AAPD)

Founded in 1947, the AAPD is a not-for-profit membership association representing the specialty of pediatric dentistry. The AAPD's 7,800 members are primary oral health care providers who offer comprehensive specialty treatment for millions of infants, children, adolescents and individuals with special health care needs. The AAPD also represents general dentists who treat a significant number of children in their practices. As advocates for children's oral health, the AAPD develops and promotes evidence-based policies and guidelines, fosters research, contributes to scholarly work concerning pediatric oral health, and educates health care providers, policymakers and the public on ways to improve children's oral health. For further information, please visit the AAPD Web site at www.aapd.org.

American Academy of Pediatric Dentistry

211 East Chicago Avenue, Suite 1700

Chicago, Illinois 60611

(312) 337-2169

aapdinfo@aapd.org

www.aapd.org



©2011 American Academy of Pediatric Dentistry
All rights reserved.